



21. Donier Grejie





THE EFFECT OF FOLIAR APPLICATION OF AGROSTEMIN® ON MUSK MELON PLANTATION

Phenological Observations –

(Cucumis melo)





These experiments were conducted in the municipality of Mossoro and Baraunas – RN, respectively, Brazil.

AGROSTEMIN® was applied by fertigation with water solution between the 5th and 7th day after transplanting, at the quantity of 30 g per hectare.



PHENOLOGICAL OBSERVATIONS

(photos)



FIRST EXPERIMENT


Sort: GALIA

Variety: ESTORIL

AGROSTEMIN® was applied: 5 days after the transplanting

AGE of plants: 21 days after transplanting and 16 days after the application











SECOND EXPERIMENT

Sort: AMARELO

Hybrid: GOLD EX

AGROSTEMIN® was applied: 7 days after the transplanting

AGE of plants: 23 days after transplanting and 16 days after the application











CONTROL

TREATED




THIRD EXPERIMENT

Sort: GALIA


Variety: DENNY

AGROSTEMIN® was applied: 7 days after the transplanting

AGE of plants: 22 days after transplanting and 15 days after the application









FOURTH EXPERIMENT

Sorta: GALIA

Varijetet: 8530

AGROSTEMIN[®] was applied: 5 days after the transplanting

AGE of plants: 20 days after transplanting and 15 days after the application















22. Donier Grejie

www.agrostemin.com