

AGROSTEMIN[®]



Dr. Danilo Gajić



**EFEKAT PRIMENE AGROSTEMIN[®]-a
PUTEM TRETIRANJA SEMENA NA KRMNI SIRAK
(*Sorghum bicolor*)
- proizvodni ogled -**



Ogled je izveden 2010., u gradu Pentecoste –CE na Oglednom imanju Vale do Curu Federalnog Univerziteta Ceara.

AGROSTEMIN[®] je apliciran na dan setve tretiranjem semena dozom proporcijalnoj standardnoj: 30 g **AGROSTEMIN**[®]-a na količinu semena za zasejavanje jednog hektara.


Efekti **AGROSTEMIN**[®]-a bili su testirani uporedno sa tri nivoa đubrenja, i to 0%, 50% i 100% preporučene doze đubriva za tu kulturu.


Tabela, grafikon i slike ilustruju stanje useva 86. dana od dana setve.


PROSEČNA TEŽINA DVA BOKORA

ĐUBRIVO			
količina	SA AGROSTEMIN®-om	BEZ Agrostemina	POVEĆANJE (%)
0%	167,79	45,67	+267
50%	170,87	74,09	+131
100%	165,20	77,64	+113

Prosečna težina dva bokora biljka tretiranih i sa **AGROSTEMIN®**-om
u poređenju sa **KONTROLOM** ("100% đubrivo").










PROSEČNA TEŽINA DVA SVEŽA KORENA

ĐUBRIVO			
količina	SA AGROSTEMIN [®] -om	BEZ Agrostemina	POVEĆANJE (%)
0%	42,76	24,53	+ 74
50%	72,79	28,37	+ 157
100%	60,97	37,05	+ 65








**0%
AGR**

**50%
AGR**

**100%
AGR**

100%







AGROSTEMIN



Dr. Danilo Gajić

www.agrostemin.com